

0755-1055, WEDNESDAY, 13 DECEMBER, 2017


NAME: _____	ALPHA NUMBER: _____
INSTRUCTOR: _____	SECTION: _____

Write your name, alpha number, section number and instructor's name on this page and on your bubble sheet. Bubble in your alpha number in the space provided on the left side of your bubble sheet.

PART ONE. Multiple Choice (50%). The first 20 problems are multiple choice. Fill in the letter for the best answer on the bubble sheet. There is no additional penalty for a wrong answer. You may use your TI-36X Pro calculator. The transformations from cartesian to cylindrical and spherical coordinates are included at the bottom of the last page of this test.

1. A contour map for a function $f(x, y)$ is shown below. Which of the following is true at the point P ?

- (a) $f_x(P) > 0$ and $f_y(P) > 0$
- (b) $f_x(P) > 0$ and $f_y(P) < 0$
- (c) $f_x(P) < 0$ and $f_y(P) > 0$
- (d) $f_x(P) < 0$ and $f_y(P) < 0$
- (e) $f_x(P) = 1$ and $f_y(P) = 0$



2. Let $f(x, y)$ be the function from problem 1. Which of the following vectors approximates best the direction of $\nabla f(P)$?

- (a) $\langle 1, 0 \rangle$
- (b) $\langle 0, 1 \rangle$
- (c) $\langle 1, -1 \rangle$
- (d) $\langle -1, 1 \rangle$
- (e) $\nabla f(P)$ is the zero vector

3. The directional derivative of the function $g(x, y, z) = z^3 - x^2y$ at the point $(1, 6, 2)$ in the direction of the vector $\langle 3, 4, 12 \rangle$ is equal to

- (a) 13
- (b) 41
- (c) 104
- (d) 0
- (e) 8

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

4. If $T(x, y) = 2x^2 + xy + y^2$ is the temperature at the point (x, y) on the plane, in which direction should one move from the point $(1, 2)$ in order for the temperature to increase the fastest?

- (a) $\langle 1, 2 \rangle$
 - (b) $\langle 2, 1 \rangle$
 - (c) $\langle 4, 4 \rangle$
 - (d) $\langle 5, 6 \rangle$
 - (e) $\langle 6, 5 \rangle$
-

5. If $z = f(x, y)$ and $x = g(t)$, $y = h(t)$ where f , g , and h are differentiable functions such that

$$g(3) = 2, \quad g'(3) = 5, \quad h(3) = -1, \quad h'(3) = 2$$

$$f_x(2, -1) = 4, \quad f_y(2, -1) = -2$$

then $\frac{dz}{dt}$ when $t = 3$ is equal to

- (a) 44
 - (b) 16
 - (c) 10
 - (d) 8
 - (e) 2
-

6. If the velocity of a particle at time t is $\vec{v}(t) = \langle 2 \sin t, \cos t \rangle$ and its position at time $t = 0$ is $\vec{r}(0) = \langle 1, -1 \rangle$ then its position $\vec{r}(\pi)$ at time $t = \pi$ is

- (a) $\langle 2, 1 \rangle$
 - (b) $\langle 3, 1 \rangle$
 - (c) $\langle 3, -1 \rangle$
 - (d) $\langle 5, 1 \rangle$
 - (e) $\langle 5, -1 \rangle$
-

7. A tangent vector to the curve $\mathbf{r}(t) = \langle t + 1, t^2, t^2 \rangle$ at the point $(0, 1, 1)$ is

- (a) $\langle 1, 2t, 2t \rangle$
- (b) $\langle 1, -2, -2 \rangle$
- (c) $\langle 1, 0, 0 \rangle$
- (d) $\langle 1, 2, 2 \rangle$
- (e) $\langle 0, 2, 2 \rangle$

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

8. The length of the parabolic arc $x = y^2$ from $(0, 0)$ to $(4, 2)$ is equal to

- (a) $\sqrt{20}$
 - (b) $\int_0^4 \sqrt{t^2 + t^4} dt$
 - (c) $\int_0^2 \sqrt{1 + 4t^2} dt$
 - (d) $\int_0^4 \int_0^2 \sqrt{x^2 + y^4} dy dx$
 - (e) $\int_0^4 \int_0^2 \sqrt{4x^2 + y} dy dx$
-

9. The tangent plane to the surface $z = 4 - x^2 + y$ at the point $(1, 2, 5)$ has equation

- (a) $-2x + y + z = 5$
 - (b) $z = 5 + 2x(x - 1) + (y - 2)$
 - (c) $2x - y + z = 0$
 - (d) $-2x + y + z = 0$
 - (e) $2x - y + z = 5$
-

10. Let $f(t, p)$ be the speed of sound, in meters per second, when the temperature is t degrees Celsius and the pressure is p atmospheres. The statement $\frac{\partial f}{\partial t}(2, 10) = 0.4$ means

- (a) When the temperature is 2 degrees Celsius and the pressure 10 atmospheres, the speed of sound is 0.4 meters per second.
- (b) When the temperature is 2 degrees Celsius and the pressure 10 atmospheres, the speed of sound increases by 40%.
- (c) When the temperature is 2 degrees Celsius and the pressure 10 atmospheres, then for every degree of increase in temperature the speed of sound increases by approximately 0.4 meters per second.
- (d) When the temperature is 2 degrees Celsius and the pressure 10 atmospheres, the speed of sound increases by approximately 0.4 meters per second.
- (e) When the temperature is 2 degrees Celsius and the pressure 10 atmospheres, the maximum rate of increase of the speed of sound is 0.4 meters per second.

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

11. Let D be the region on the plane between the circles $x^2 + y^2 = 4$ and $x^2 + y^2 = 1$. Which of the following is equal to $\iint_D 1 \, dA$?

(I) $\int_0^{2\pi} \int_1^2 r \, dr \, d\theta$ (II) $2 \int_{-2}^2 (\sqrt{4-x^2} - \sqrt{1-x^2}) \, dx$ (III) $\iint_D (2x - 2y) \, dA$

- (a) (I) only
 (b) (II) only
 (c) (I) and (II) only
 (d) (I) and (III) only
 (e) (I), (II) and (III)

12. The volume of the solid region inside the sphere $x^2 + y^2 + z^2 = 4$, underneath the cone $z = \sqrt{x^2 + y^2}$ and above the xy -plane is given by

(a) $\int_0^{2\pi} \int_0^2 \int_0^r r \, dz \, dr \, d\theta$
 (b) $\int_0^{2\pi} \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \int_0^2 \rho^2 \sin \phi \, d\rho \, d\phi \, d\theta$
 (c) $\int_0^{2\pi} \int_0^2 \int_r^{\sqrt{4-r^2}} 1 \, dz \, dr \, d\theta$
 (d) $\int_0^{2\pi} \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \int_0^2 1 \, d\rho \, d\phi \, d\theta$
 (e) $\int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} (\sqrt{x^2 + y^2} - \sqrt{4 - x^2 - y^2}) \, dx \, dy$

13. Reversing the order of integration, the integral $\int_0^1 \int_{x^2}^{\sqrt{x}} xy^2 \, dy \, dx$ becomes

(a) $\int_0^1 \int_{\sqrt{y}}^{y^2} xy^2 \, dx \, dy$
 (b) $\int_0^1 \int_{y^2}^{\sqrt{y}} xy^2 \, dx \, dy$
 (c) $\int_0^1 \int_{\sqrt{y}}^{y^2} yx^2 \, dx \, dy$
 (d) $\int_{x^2}^{\sqrt{x}} \int_0^1 xy^2 \, dx \, dy$
 (e) $\int_{\sqrt{x}}^{x^2} \int_0^1 xy^2 \, dx \, dy$

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

14. Which of the following is a parametrization for the cone $y^2 = x^2 + z^2$?

- (a) $\mathbf{r}(u, v) = \langle \cos v, u, \sin v \rangle$
 - (b) $\mathbf{r}(u, v) = \langle \cos v, v, \sin v \rangle$
 - (c) $\mathbf{r}(u, v) = \langle u \cos v, 1, u \sin v \rangle$
 - (d) $\mathbf{r}(u, v) = \langle u \cos v, u, u \sin v \rangle$
 - (e) $\mathbf{r}(u, v) = \langle \cos v, 1, \sin v \rangle$
-

15. The work done by the force field $\mathbf{F}(x, y) = \langle y, x \rangle$ along the straight line segment from the point $(1, 1)$ to the point $(2, 4)$ is equal to

- (a) 0
 - (b) 7
 - (c) 13
 - (d) 14
 - (e) 28
-

16. For which value of the constant k is the vector field $\mathbf{F}(x, y) = \langle y^2 + ky \sin(2x), 2xy + \cos(2x) \rangle$ conservative?

- (a) -2
 - (b) -1
 - (c) 0
 - (d) 1
 - (e) 2
-

17. Suppose that $\mathbf{F}(x, y) = \langle P(x, y), Q(x, y) \rangle$ is a vector field whose components have continuous partial derivatives in \mathbb{R}^2 . Let D be the disk $x^2 + y^2 \leq 1$ and C its boundary. If it is known that $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$, which of the following must be true?

(I) \mathbf{F} is conservative

(II) $\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} = 0$

(III) $\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA = 0.$

- (a) (I) only
- (b) (II) only
- (c) (III) only
- (d) (I), (II) and (III)
- (e) None of the three statements is necessarily true

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

18. The outward flux $\iint_S \mathbf{F} \cdot d\mathbf{S}$ of the vector field $\mathbf{F}(x, y, z) = \langle 3x - 4xz, y^2 + xz, 2z^2 - 2yz \rangle$ through the surface S which is the boundary of the solid cube $0 \leq x \leq 2$, $0 \leq y \leq 2$, $0 \leq z \leq 2$ is

- (a) 24
 - (b) 0
 - (c) 3
 - (d) 8
 - (e) -8
-

19. Suppose that \mathbf{F} is a vector field whose components have continuous partial derivatives in \mathbb{R}^3 . Let S be the sphere $x^2 + y^2 + z^2 = 1$ with the outward orientation, S_1 the top hemisphere with the upward orientation and S_2 the bottom hemisphere with the downward orientation. If it is known that $\text{div}\mathbf{F} = 0$, which of the following statements must be true?

(I) \mathbf{F} is conservative.

(II) $\iint_S \mathbf{F} \cdot d\mathbf{S} = 0$.

(III) $\iint_{S_1} \mathbf{F} \cdot d\mathbf{S} = \iint_{S_2} \mathbf{F} \cdot d\mathbf{S}$.

- (a) (I) only
 - (b) (II) only
 - (c) (I) and (II) only
 - (d) (I) and (III) only
 - (e) (I), (II) and (III)
-

20. If C is the curve consisting of the semicircle $x^2 + y^2 = 4$, $y \geq 0$ from $(2, 0)$ to $(-2, 0)$ followed by the straight line segment from $(-2, 0)$ to $(2, 0)$, then the line integral

$\int_C (x^2 - y) dx + (x + y^3) dy$ is equal to

- (a) -8π
- (b) -4π
- (c) 0
- (d) 4π
- (e) 8π

END OF MULTIPLE CHOICE PORTION OF THE TEST

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

PART TWO. Free response (50%). The remaining 10 problems are not multiple choice. Each is worth 10 points. Answer them in the space provided on these pages. Show all your work. You may use your TI-36X Pro calculator. No other aides are allowed.

21. (10 points) The table below shows values of the function $f(x, y)$, defined on the rectangle $R = [0, 8] \times [0, 20]$.

(a) Estimate the values of $f_x(4, 10)$ and $f_y(4, 10)$.

$y \setminus x$	0	2	4	6	8
0	20	19	13	11	8
5	23	20	14	13	11
10	28	25	20	17	14
15	33	31	24	19	14
20	41	40	30	26	20


(b) Use linear approximation to estimate $f(4.2, 9.9)$.

(c) Estimate $\iint_R f(x, y) dA$ using the midpoint rule with four sub-rectangles of equal size.

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

22. (10 points) Consider the function $f(x, y) = x^3 - y$.

(a) Sketch a contour map of f that includes the level curves at levels $z = 0, -1$ and -2 .



(b) On the contour map from part (a) and at the point $(1, 2)$, sketch accurately the direction in which f increases at the fastest possible rate.

(c) What is the maximum rate of increase of f at the point $(1, 2)$?

23. (10 points) Find the critical points of the function $f(x, y) = xy^2 - x + 2y$ and determine whether they are points of local maximum, local minimum or saddle points.

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

24. (10 points) Sketch the region E bounded by the paraboloid $z = 9 - x^2 - y^2$ and the plane $z = 5$ and compute its volume.

25. (10 points) Let W be the solid region in the first octant bounded by the plane $4x + 2y + z = 8$ and the three coordinate planes. If the density at the point (x, y, z) is numerically equal to the distance from the xy -plane, set up a triple iterated integral equal to the mass of W . Do not evaluate the integral.

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

26. (10 points) Consider the vector field $\mathbf{F}(x, y, z) = \langle 2xy + z + 3x^2, x^2 + 2yz, x + y^2 \rangle$.

(a) Show that \mathbf{F} is conservative and find a potential function.

(b) Compute $\int_C \mathbf{F} \cdot d\mathbf{r}$ where C is the curve $\mathbf{r}(t) = \langle \cos t, \sin t, 2 - \sin t \rangle$, $0 \leq t \leq \pi$.

27. (10 points) Compute the flux $\iint_S \mathbf{F} \cdot d\mathbf{S}$ of the vector field $\mathbf{F}(x, y, z) = \langle x, y, z \rangle$ through the cylinder $x^2 + y^2 = 4$, $0 \leq z \leq 3$ with the outward orientation (i.e., the normal vector points away from the z -axis).

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

28. (10 points) Let S be the part of the sphere $x^2 + y^2 + z^2 = 8$ that lies above the plane $z = 2$. Set up a double iterated integral that is equal to the surface area of S . Do not evaluate the integral.

29. (10 points) Compute the surface integral $\iint_S \text{curl} \mathbf{F} \cdot d\mathbf{S}$ where $\mathbf{F}(x, y, z) = \langle z + y, x^2z, -x + z^2y \rangle$ and S is the paraboloid $y = 1 - x^2 - z^2$, $y \geq 0$ with the outward orientation (i.e., the normal vector has positive y component).

0755-1055, WEDNESDAY, 13 DECEMBER, 2017

30. (10 points) Prove the following Theorem: Let C be a smooth curve given by the vector function $\mathbf{r}(t)$, $a \leq t \leq b$. Let f be a differentiable function of three variables whose gradient vector ∇f is continuous on C . Then

$$\int_C \nabla f \cdot d\mathbf{r} = f(\mathbf{r}(b)) - f(\mathbf{r}(a))$$

Spherical Coordinates

$$\begin{aligned}x &= \rho \sin \phi \cos \theta \\y &= \rho \sin \phi \sin \theta \\z &= \rho \cos \phi\end{aligned}$$

Cylindrical Coordinates

$$\begin{aligned}x &= r \cos \theta \\y &= r \sin \theta \\z &= z\end{aligned}$$