

## Lesson 3. Graphical Solution of Optimization Models

### 0 Warm up

On the axes on page 2, draw the following equations, and label the points of intersection.

$$4C + 2V = 32 \quad 4C + 6V = 48$$

### 1 Overview

- Last time, we formulated a linear program for Farmer Jones's problem:

$C$  = number of chocolate cakes to bake


$V$  = number of vanilla cakes to bake

$$\begin{aligned} \text{maximize} \quad & 3C + 4V \\ \text{subject to} \quad & 4C + 2V \leq 32 & (1) \\ & 4C + 6V \leq 48 & (2) \\ & C \geq 0 & (3) \\ & V \geq 0 & (4) \end{aligned}$$

- By trial-and-error, the best feasible solution we found was  $C = 6$ ,  $V = 4$
- Today, we will discuss how to solve Farmer Jones's model systematically

## 2 Solving Farmer Jones's model graphically


- We can graphically solve linear programs with 2 variables
- The feasible region – the collection of all feasible solutions – for Farmer Jones's optimization model:



- Any point in this shaded region represents a feasible solution
- How do we find the one with the highest value?
- $C = 2, V = 2$  is a feasible solution with value
- The set of values of  $C$  and  $V$  with the same value satisfies:
- Idea:
  - Draw lines of the form  $3C + 4V = k$  for different values of  $k$
  - Find the largest value of  $k$  such that the line  $3C + 4V = k$  intersects the feasible region
- These lines are called **contour plots**
  - Lines through points having equal objective function value

### 3 Sensitivity analysis

- For what profit margins on vanilla cakes will the current optimal solution remain optimal?



- Key observation:

- Slope of (1) = , slope of (2) =


- Let  $a$  be the new profit margin on vanilla cakes

⇒ objective function is , slope of contour plots =


⇒ If , then the current optimal solution remains optimal

## 4 Outcomes of optimization models


- An optimization model may:
  1. have a **unique optimal solution**
    - e.g. the original Farmer Jones model
  2. have **multiple optimal solutions**
    - e.g. What if the profit margin on chocolate and vanilla cakes is \$2 and \$3, respectively, instead?
    - Farmer Jones's objective function is then



3. be **infeasible**: no choice of decision variables satisfies all constraints
  - e.g. What if the demands of Farmer Jones's neighbors dictate that he needs to bake at least 9 chocolate cakes?
  - Then we need to add the constraint



4. be **unbounded**: for any feasible solution, there exists another feasible solution with a better value
- e.g. What if the circumstances have changed so that the feasible region of Farmer Jones's model actually looks like this:



## 5 Next time...

- More linear programming models
- Introduction to GMPL (bring your laptops)